

# Prevention of adhesion bands by ibuprofen-loaded PLGA nanofibers

*Fatemeh Jamshidi-Adegani<sup>1, 2</sup>, Ehsan Seyedjafari<sup>3</sup>, Nematollah Gheibi<sup>4</sup>, Masoud Soleimani<sup>2, 5\*</sup>, Mehdi Sahmani<sup>6\*\*</sup>*

<sup>1</sup>Department of Molecular Medicine, School of Medicine, Qazvin University of Medical Science, Qazvin, Iran

<sup>2</sup>Department of Molecular Biology and Genetic Engineering, Stem Cell Technology Research Center, Tehran, Iran

<sup>3</sup>Department of Biotechnology, College of Science, University of Tehran, Tehran, Iran

<sup>4</sup>Department of Physiology and Medical Physics, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>5</sup>Department of Hematology, Faculty of Medical Science, Tarbiat Modares University, Tehran, Iran

<sup>6</sup>Department of Clinical Biochemistry and Genetics, Cellular and Molecular Research Center, Faculty of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

**\*\*Corresponding author:** Mehdi Sahmani, Department of Clinical Biochemistry, Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran. E-mail: [m.sahmani@gmail.com](mailto:m.sahmani@gmail.com)

**\*Co-Corresponding author:** MasoudSoleimani, Department of Hematology, Faculty of Medical Science, TarbiatModares University, Tehran, Iran P.O. Box 14115-111; E-mail: [soleim\\_m@modares.ac.ir](mailto:soleim_m@modares.ac.ir)

**This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/btpr.2270**

**© 2016 American Institute of Chemical Engineers Biotechnol Prog**

**Received: Sep 01, 2015; Revised: Mar 31, 2016; Accepted: Apr 01, 2016**

**Abstract**

In this study, prevention of the adhesion bands and inflammatory features has been investigated using poly (lactic-co-glycolic acid)-ibuprofen (PLGA-IB) nanofibrous meshes in a mice model. In order to find the optimized membrane for prevention of postoperative adhesion bands, we have compared PLGA-IB group with PLGA, IB and control groups in a mice adhesion model. Two scoring adhesion systems were used to represent the outcome. According to the results obtained in this study, the PLGA-IB nanofiber membrane showed a greater reduction in adhesion band than other groups. In conclusion, among FDA-approved polymers and drugs, PLGA-IB meshes could be applicable as a potential candidate for prevention of post-operative abdominal inflammation and adhesion bands formation.

**Key Words:** intra-peritoneal adhesion; poly (lactic-co-glycolic acid) (PLGA); ibuprofen